

CHEMISTRY

** This course meets the 1 hr. Chemistry requirement for Florida license renewal. **

THYROID HORMONE EXCESS: GRAVES DISEASE

COURSE CODE: C021
CONTACT HOURS: 3
COURSE LEVEL: Intermediate
CE BROKER #: 20-404921

-- COURSES ARE REVIEWED EVERY 2 YEARS --

Continuing Education Unlimited

6231 PGA Blvd / Suite 104, #306 / Palm Beach Gardens, FL 33418

888-423-8462 / General Fax: 561-775-4933 / Answer Sheet ONLY Fax: 561-775-4948 / www.4CEUINC.com

LAB PROVIDER #s: Florida: 50-2256 | ASCLS P.A.C.E.: 511 | CALIFORNIA ACCREDITING AGENCY (AA) #: 0001

COURSE OBJECTIVES

1. Recall the clinical features of Graves' disease.
2. Recall the background of Graves' disease including the epidemiology and risk factors.
3. Discuss the pathogenesis of Graves' disease.
4. Describe the pathogenic mechanisms associated with Graves' disease.
5. List the clinical features associated with Graves' disease.
6. Recall the tests available to diagnose Graves' disease.
7. List the treatment options available for Graves' disease.

COPYRIGHT INFO:

AUTHOR / INFORMATION SOURCE

Licensors/Publisher: InTech Open

Text: Thyroid Hormone, Chapter

6 Author(s): N. Agarwal, et al

R : October 2019

PERMISSIONS

This is an Open Access article distributed under the terms of the Creative Commons Attribution License 3.0 (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted Sharing (use, distribution, and reproduction), Remixing (adapting the work), or making commercial use of the work provided that the original work, author, and licensor are properly cited. Use of this article in no way suggests endorsement of our company by the authors or licensor.

Access article directly:

<http://www.intechopen.com/books/thyroid-hormone/thyroid-hormone-excess-graves-disease>

ARTICLE REPRODUCTION

Any article reproduction falls under the guidelines of the Creative Commons License listed above.

Copyright is retained by the original author(s) and proper, citation must be kept intact. Reproduction requires that integrity is maintained and its original authors, citation details, and publisher are identified.

CEUINC LICENSE INFO:

LABORATORY

CA Department of Health:	0001
Florida Board of Clinical Lab:	50-2256
ASCLS P.A.C.E.®	511

Our courses are accepted by: AMTIE, AMT, ASCP, CA, FL, LA, ND, NV, MT, RI, TN, WV

** If you do not see your organization, state, or licensing agency listed above it does not mean that the credits will be unacceptable. Most licensing bodies accept **P.A.C.E.** credits, so please check directly with them for acceptance of our courses.**

PHLEBOTOMY

Most licensing bodies accept **P.A.C.E.** credits. Please check with them directly for acceptance of our course credits.

OTHER MEDICAL DISCIPLINES

Many medical licensing bodies will accept credits issued by valid licensed providers of other disciplines. Please check directly with the state, agency, or organization that issued your license for acceptance of our credits.

** CEUINC is approved as a provider of continuing education programs in the clinical laboratory sciences by the ASCLS P.A.C.E.® Program. **



Continuing Education Unlimited

6231 PGA Blvd, Suite 104 / #306 / Palm Beach Gardens, FL 33418
Phone: 888-423-8462 / Fax: 561-775-4933 / Web: www.4CEUINC.com / Email: Lab@4ceuinc.com

Thanks for choosing 4CEUINC for your continuing education needs! We strive to offer you current, cost effective course material. If you have comments or suggestions, be sure to add them to your evaluation. If you like our courses pass them on to a coworker or friend.

READ BEFORE COMPLETING MATERIAL



Last Revised 04/14/19

GENERAL COURSE INFO:

- 1.) Check your courses to make sure they are the correct titles that you ordered.
- 2.) All quiz answers can be found within the reading material and typically follow in order of the reading.
- 3.) All courses are reviewed every 2 yrs, at which time they will either be expired from our course list, updated, or renewed for an additional 2 years. Expiration dates can be found on this course list: https://4ceuinc.com/site/assets/pdf/current_course_list.pdf
- 4.) All of your records are available to you online through our website, even for Home Study. If you don't have a login to access your records, please contact our office and we will give you that information. **Please do not create a 2nd profile!**

HOME STUDY COURSES: (Combo Courses only)

- 1.) If you have multiple answer sheets, please make sure you're using the correct one.
- 2.) **CHANGES:** Home Study Complete formats have been replaced by "Book Only" formats, where you can purchase the printed book, then add on the quiz format option of your choice - - "Online Quiz Only" or "Answer Sheet Only".
- 3.) **CHANGES:** Answer sheets will no longer be mailed. You will print them out by logging into your online profile and visiting the My Courses list. Locate the correct course, then print your answer sheet. You may reprint as needed.
- 4.) **CHANGES:** As of May 1, 2019, answer sheets may **no longer** be returned by mail. You may submit them via fax to 561-775-4948 **OR** upload an image, doc, or pdf file by visiting your My Courses list online. As an alternate option, all Home Study quizzes **can be completed online**, which allows for **immediate** grading and certificate printing.
- 5.) **CHANGES:** As of May 1, 2019, we will no longer be returning Certificates of Completion. They may be printed out any time by visiting your "Course History" in your online profile. In the right column, take the Eval, then print the Certificate.

ONLINE COURSES:

- 1.) Course materials are currently offered in Adobe PDF documents, PowerPoint (PPT), and Video.
- 2.) You can read PDF & PPT documents online or print them out for reference. You may also store it on your hard drive or copy it to flash drive. Per copyright law, you're entitled to make **one copy** of each online course.
- 3.) Quizzes can be printed out if you'd like to work offline by clicking <Print Copy of Quiz>. Once you are done, simply transfer your answers to the active, online <Take Quiz> link and then click "Grade".
- 4.) When completing the online quiz, **it's important** to enter the answers without interruption, as some of the newer anti-virus programs may log you out of active content after a short period of time, causing you to lose your answers.

REGISTERED GROUPS:

- 1.) It's important to keep the Educational Coordinator's contact information current with our office at all times.
- 2.) In order to receive the group discount, you must be listed as a group member **OR** have the registration code to add to your online profile.
- 3.) You can share reading material with your co-workers for Combo Courses or Textbooks. They can purchase just the **online quiz** for that course allowing them to immediately complete the quiz and print their certificate of completion.
- 4.) **Have 5 or more employees?** Set up a Group Account & receive discounts on continuing education for your staff.



ALERT: When faxing, make sure to include a cover sheet with your full name & phone #. When faxing or uploading your Home Study answer sheets, make a note of the date and time you sent them! This safeguards you in the event that your answer sheet does not transmit properly to 4CEUINC.



Continuing Education Unlimited

6231 PGA Blvd, Suite 104 / #306 / Palm Beach Gardens, FL 33418
Phone: 888-423-8462 / Fax: 561-775-4933 / Web: www.4CEUINC.com / Email: Lab@4ceuinc.com



FREQUENTLY ASKED QUESTIONS

Last Revised 04/14/19

Q. Your phones are often busy, how can I reach you?

A. Because we have a small staff, the most efficient way would be to fill out a “Contact Us” form. On the main menu at www.4CEUINC.com, go to ‘Support’, ‘Contact Us’, then fill out the form. Be sure to enter your full information.

Q. I ordered a Home Study Combo Course, may I complete the quiz online?

A. Yes, quizzes from all course formats can be completed online, just log into your profile and go to “My Courses”.

Q. I need my certificate dated on a certain date; how can I be sure that this will happen?

A. 1.) Complete the quiz online. **OR** 2.) Be sure to fax or upload your answer sheet at least 72 hours before that date.

Q. What score is considered passing?

A. A score of 70% or higher is considered a passing grade. You may repeat the quiz if you fail it the first time.

Q. How can I return my Home Study answer sheet to your office?

A. CHANGES: As of May 15, 2019, answer sheets may no longer be returned by mail. They may **ONLY** be returned by fax to **561-775-4948** **OR** uploaded directly into our system through your My Courses list in your online profile. Be sure to keep a record of the date & time you send the Answer Sheet in the event that there is a transmission problem!

Q. May I share my Combo Course or Textbook with others?

A. YES, Combo Courses and Textbooks can be shared. One person will buy the “complete course” package & each additional person will purchase an “online quiz only”. Remember to share **both** the reading material **and** the quiz.

Q. How will I get my Certificate of Completion for a Home Study Combo Course?

A. CHANGES: As of May 1st, 2019, we will no longer be returning CEU certificates by mail due to past mail delivery problems. You may print your Certificate any time by logging into your online profile and visiting your course history.

Q. May I print out an online course?

A. YES, you may print **1 copy** of an online course. Copyright laws do not allow more than one copy to be printed!

Q. Where can I find additional information out about a course?

A. Basic course information is located on the course cover. Additional information is located on the 2nd page of each course.

Q. What are your most popular courses?

A. Currently our most popular courses remain the 12 Month Online Subscription and our Combo Courses.

Q. Does CEUINC offer group discounts or group packages?

A. YES. We require at least 5 participants and a person to act as the educational coordinator for the group. Discounted pricing depends on the number of participants and the package chosen. Please call 561-775-4944 Ext. 4 for details.

Q. Does CEUINC report to CE Broker for Florida licensees?

A. YES. Once you complete the online quiz, your course completions are sent to CE Broker immediately, provided that we have your correct Florida license number listed in your profile. Allow 48 hours for faxed or uploaded answer sheets.



ALERT: When faxing, make sure to include a cover sheet with your full name & phone #. When faxing or uploading your Home Study answer sheets, make a note of the date and time you sent them! This safeguards you in the event that your answer sheet does not transmit properly to 4CEUINC.



THYROID HORMONE EXCESS: GRAVES DISEASE

Category: Chemistry | **Contact Hours:** 3 | **Course Code:** C021

- 1.) Patients presenting with toxic features can be because of thyroiditis which typically is a few weeks duration or Graves' disease, which is for several months duration most of the time.
 - A. True
 - B. False

- 2.) Graves' disease is the most common cause of thyrotoxicosis and it accounts for _____ of the cases of thyrotoxicosis.
 - A. 25%
 - B. 40-50%
 - C. 60-80%

- 3.) The postpartum period is associated with a fourfold to eightfold increased risk for the onset of Graves' disease.
 - A. True
 - B. False

- 4.) Thyroid stimulating antibodies (TSAb) are produced mainly by the _____ infiltrating the thyroid gland and _____ present in the draining lymph nodes, and partly by the circulating blood _____.
 - A. monocytes
 - B. neutrophils
 - C. lymphocytes

- 5.) TH1 cells are mainly involved in delayed type of hypersensitivity reactions, and it produces the cytokines like TNF-B, IFN-Y, IL-2, IL-10 and IL-17.
 - A. True
 - B. False

- 6.) After treatment with antithyroid drugs and radioiodine, the vascularity of the thyroid gland _____, follicles enlarges and filled with colloid, and papillary projection regresses.
 - A. increases
 - B. decreases
 - C. remains unchanged

- 7.) Clinical ophthalmopathy is present in about 50% of Graves' disease patients.
 - A. 20%
 - B. 37%
 - C. 50%

- 8.) The typical lesion is an inflamed, indurated plaque with a deep pink or purple color and an orange skin appearance.
 - A. True
 - B. False

**THYROID HORMONE EXCESS: GRAVES DISEASE
- QUIZ PAGE 2 -**

- 9.) Serum TSH level is suppressed or undetectable with _____ thyroxine (TT4) and triiodothyroxine (TT3) levels in patients of Graves' disease.
A. increased
B. decreased
C. normal
- 10.) Anti thyroid peroxidase (TPO) antibody can be detected in up to ___ of patients with Graves' disease whereas anti-thyroglobulin antibody is present in _____ cases.
A. 40%, 30-70%
B. 65%, 80-90%
C. 90%, 50-80%
- 11.) Dexamethasone in high doses (8mg/day) decreases the T4 secretion by the thyroid gland, inhibit the peripheral conversion of T4 to T3, and has immunosuppressive effect.
A. True
B. False
- 12.) Radioiodine is one of the first line therapies for the Graves' disease, with _____ as the agent of choice.
A. I51
B. I131
C. I236
- 13.) Antithyroid drugs should be stopped _____ prior to radioiodine treatment and should be restarted after _____ when required.
A. 10-14 days, 11 months
B. 3 to 8 days, 7 days
C. 2 months, 14 days
- 14.) Hypoparathyroidism is a possible complication to thyroid surgery.
A. True
B. False
- 15.) Thyroid storm or thyrotoxic crisis is a life threatening exacerbation of hyperthyroidism.
A. True
B. False

*******LAST QUESTION *******

Thyroid Hormone Excess: Graves' Disease

N.K. Agrawal, Ved Prakash and Manuj Sharma

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/47864>

1. Introduction

Thyroid hormone excess due to any cause is state of thyrotoxicosis, whereas hyperthyroidism is a state of thyroid hormone excess due to hyperfunctioning of thyroid gland. The major causes of thyrotoxicosis include Graves' disease, solitary toxic adenoma, and toxic multinodular goiter. In thyrotoxicosis, free hormone levels are invariably increased. The reverse is not true in that increased free thyroid hormone levels do not always point to thyrotoxicosis. In illness or resistance to thyroid hormones increased free hormone levels are present while the patients are clinically euthyroid or even sometimes hypothyroid. Subclinical thyrotoxicosis is defined as a state in which free thyroid hormones (FT4 & FT3) are within normal limit, but serum TSH level is low. The most common cause of subclinical thyrotoxicosis is exogenous administration of thyroid hormone rather than Graves' disease. Patients of subclinical thyrotoxicosis either have no symptoms or have mild non specific symptoms.

The various causes of thyrotoxicosis are listed below in table 1

1.1. Clinical features

Clinical manifestations of the thyrotoxicosis are similar for various causes of thyrotoxicosis. However certain features provide some clues about specific cause of throtoxicosis. These features include the duration and mode of onset of thyrotoxicosis, size and shape of the thyroid gland, presence or absence of the extra-thyroidal manifestations like Graves' eye sign, pre-tibial myxoedema, acropachy. Patient presenting with toxic features can be because of thyroiditis or Graves' disease, but the symptoms are of few weeks duration in former while in later condition, it is for several months, most of the time. The common clinical features of thyrotoxicosis are listed below in table 2.

<p>A. Thyroid hormone excess with hyperthyroidism</p> <ol style="list-style-type: none"> a. Primary Hyperthyroidism <ol style="list-style-type: none"> i. Graves' Disease ii. Toxic Thyroid adenoma iii. Toxic Multinodular Goiter iv. Metastases from Thyroid carcinoma v. Mutations of TSH receptor, G_{α} (Mc Cune Albright Syndrome) vi. Struma Ovarii vii. Iodine ingestion (jod-Basedow phenomenon) b. Secondary hyperthyroidism <ol style="list-style-type: none"> i. TSH secreting pituitary tumours ii. Pituitary Thyroid hormone resistance <p>B. Thyroid hormone excess without hyperthyroidism</p> <ol style="list-style-type: none"> a. Throiditis: subacute, silent b. Ingestion of thyroid tissue, thyroid hormone c. Thyroid gland destruction by amiodarone, radiotherapy, infarction in thyroid adenoma

Table 1. Causes of thyroid hormone excess

Symptoms	Signs
Irritability, Hyperactivity, Dysphoria	Tachycardia, Lone Atrial Fibrillation
Heat Intolerance	Fine tremor
Palpitation, Nervousness	Warm moist skin, Goiter
Fatigue, lethargy	Proximal Myopathy
Dyspnoea	Hyperreflexia
Weight loss despite polyphagia	Lid lag, Lid tremors
Hyperdefecation, Polyuria	Gynaecomastia
Oligomenorrhoea	Loss of libido

Table 2. Common clinical features

2. Graves' disease

Diffuse toxic goitre is commonly known as Graves' disease. Classically it is characterized by hyperthyroidism and diffuse goiter. Graves' disease may be associated with an infiltrative orbitopathy and ophthalmopathy and less commonly with an infiltrative dermopathy. Infiltrative orbitopathy is a unique manifestation of Graves' disease.

2.1. Historical perspective

Robert James Graves, an Irish physician, first described 3 females with this disease in 1835¹ and for his contribution, this disease is known as Graves' disease in most parts of the world. Although a similar syndrome was described by Caleb Hiller Parry, a physician from Bath,

England, in 1825.² Carl A. Von Basedow from Germany first described the triad of exophthalmos, goiter and palpitation.³ In most non English speaking European countries the disease is still known as Basedow's disease.

2.2. Epidemiology

Graves' disease is the most common cause of thyrotoxicosis and it accounts for 60-80% of cases of thyrotoxicosis. Prevalence of Graves' disease varies with the degree of iodine sufficiency, and it is the most common cause of thyrotoxicosis in iodine sufficient countries.⁴ High dietary iodine intake is associated with an increased prevalence of Graves' disease. Prevalence of Graves' disease is about 0.4% in USA,⁵ 0.6% in Italy,⁶ and 1.1% in UK.⁷ A recent meta-analysis of various studies showed that the prevalence of Graves' disease is about 1% in general population.⁷ Prevalence of Graves' disease is 1-2% in women, and it is about 10 fold more prevalent in women than men. Peak age of onset for Graves' disease is typically in the third to fifth decade of life,⁸ but it can occur in children and in the elderly.

2.3. Risk Factors for Graves' disease

Graves' disease is a multifactorial disease in which genetic ,environmental, and hormonal factors play their role.

2.3.1. Genetic factors for Graves' disease

High prevalence of Graves' disease in family members and relatives of Graves' disease and Hashimoto's thyroiditis support that genetic factors are involved in causation of Graves' disease. There is also evidence that occurrence rate of Graves' disease is higher in monozygotic twins than dizygotic twins. The concordance rate in monozygotic twins is only 17-35% which indicate low penetrance of genes.

Graves' disease is a polygenic disease. Polymorphism in HLA-DR, CTLA-4 and PTPN-22 genes are associated with increased risk of Graves' disease. HLA-DR3 (HLA-DRB1*03), HLA-DQA1*0501 and HLA-B8 gives a risk ratio of 3-fold to 4-fold in the caucasian population. HLA-DQ3 is involved in patients with African descent whereas HLA-BW46 is involved in patients with Asian descent. The other genes involved in the pathogenesis of Graves' disease are the CD40 gene, thyroglobulin gene, TSHR gene, immunoglobulin genes, GD-1 gene (on chromosome 14q13), GD-2 gene (on chromosome 20) and GD-3 gene (on chromosome Xq21-22).

2.3.2. Environmental Factors for Graves' disease

Infection

From very early it has been suggested that Graves' disease is associated with infectious agents, but this hypothesis has not been confirmed. Incidence of recent viral infections are high in patients with Graves' disease. The association of Graves' disease with infectious

agents can be explained by molecular mimicry. Molecular mimicry implies structural similarity between infectious agents with a self antigen. Circulating antibody against *Yersinia enterocolitica* has been found in high percentage of patients with Graves' disease. Furthermore, serum from some patients recovering from *Yersinia* infections block the binding of TSH to its receptor. Low affinity binding sites for TSH have also been found with *Leishmania* and *Mycoplasma* species. However, no studies proved that infections agents have causative role in Graves' disease.

Stress

In different studies, it has been found that stressful life events precede the onset of Graves' disease. Severe emotional and physical stress, like separation from the loved one or following road traffic accident, cause release of cortisol and corticotrophin releasing hormone. So, stress is a relatively immune suppression state. Immune system overcompensates once stress is over which can precipitate disease similar to a postpartum period. In conclusion there is limited but significant evidence that stressful life events can precipitate the onset of Graves' disease in genetically susceptible individuals.

Gender

Typically Graves' disease is more prevalent in females than males. It is about 5-10 times more common in females at any age.^{6,7} In children this difference is smaller. The exact cause for female preponderance is not known, but it is similar to other autoimmune disorders. In experimental animal models of autoimmune thyroiditis it has been seen that androgens appear to down regulate the immune system.^{8,9} Other possible explanation for female preponderance is female sex steroids. But Graves' disease also occurs in men and postmenopausal women. These observations have suggested that it is the X-chromosome, not the sex steroids, which is responsible. But most of the x-linked disorders are only present in man, it has been thought that a gene with dose dependent effect on X-chromosome is responsible.

Pregnancy

Postpartum period is an important risk factor for both the onset and relapse of Graves' disease. Postpartum period is associated with a fourfold to eightfold increased risk for the onset of Graves' disease. Rebound immunity is the likely explanation for this increased risk. Graves' disease is associated with low pregnancy rate because thyrotoxicosis decreases the fertility rate. However in women with Graves' disease who became pregnant, successful pregnancy outcome is low because Graves' disease causes increased pregnancy loss and its complications. Graves' disease exacerbates during the first trimester of pregnancy and postpartum period, while it improves during the second and third trimester of pregnancy.

Smoking

Smoking is a minor risk factor for Graves' disease; however it is a major risk factor for Graves' ophthalmopathy. There are number of studies showing relationship between Graves' disease, Graves' ophthalmopathy and smoking.^{11,12}

Other risk factors:

Direct trauma to the thyroid gland, ethanol injection for the treatment of autonomously functioning thyroid nodules, or thyroid injury following radio-iodine treatment for toxic adenoma or toxic multinodular goiter are associated with an increased risk of Graves' disease. Radio-iodine treatment may also cause onset or worsening of ophthalmopathy. Possible explanation is that thyroid injury by any means cause massive release of thyroid antigens, which in turn stimulate an autoimmune reaction to TSHR in susceptible individuals.

Graves' disease onset and recurrence is also associated with iodine and iodine containing drugs like amiodarone and radio-contrast media especially in iodine deficient population.

3. Pathogenesis

Graves' disease is an example of organ specific autoimmune disorder in which both humoral and cell mediated immunity directed against different thyroid antigens are involved. TSHR is the primary autoantigen of Graves' disease, while other autoantigens like thyroglobulin and thyroid peroxidase (TPO) are secondarily involved.

3.1. Humoral immunity

TSHR is a member of GPCR super family and involves cAMP and phosphoinositol pathways for signal transduction. TSHR has large intracellular domain (subgroup B). It is a glycoprotein consisting of 744 amino acids and having molecular weight of 84 kd. A gene for TSHR is located on chromosome 14q31 and is formed by 10 exons. Circulating autoantibody directed against the TSHR is the primary factor responsible for Graves' disease. TSHR antibodies (TSHR-Ab) are of three types- namely stimulating antibody, blocking antibody and neutral antibody. Stimulating antibodies are those who after binding to TSHR activate adenylate cyclase and cause increased thyroid growth and vascularity, and increases the production of thyroid hormones. Blocking antibodies are those who after binding to TSHR act as an antagonist, whereas neutral antibodies does not have any functional activity.

Almost 50 years ago (in 1956) long acting thyroid stimulators (LATS) were discovered by Adams and Purves during a search for thyroid stimulating activity in patients with Graves' disease.¹³ Later on it was found that LATS are nothing but immunoglobulin of IgG1 subclass.

Transplacental transfer of TSHR stimulating antibodies (TSAb) from TSAb positive pregnant mother to fetus causes transient neonatal thyrotoxicosis that improves spontaneously after the disappearance of TSAb.¹⁴ This provides the definite role of TSAb in the causation of Graves' disease. TSAb are oligoclonal.

TSAb are produced mainly by the lymphocytes infiltrating the thyroid gland and lymphocytes present in the draining lymph nodes, and partly by the circulating blood lymphocytes.^{15,16} There is a positive correlation that exists between the TSAb level and serum

triiodothyronine level, serum thyroglobulin level and goiter size. TSAb are found in 90-100% of untreated Graves' disease patients.^{17, 18} Level of TSAb decreases after treatment with anti-thyroid drugs and radio-iodine.^{19, 20}

3.2. Assays for TSHR-Ab

Two types of assays are used for TSHR-Ab-Radioreceptor assays and invitro bioassays.

3.2.1. Radioreceptor assay

Radioreceptor assay is most readily available and most widely used in clinical practice. Basic principles of a radioreceptor assay is displacement of labeled TSH from solubilized TSHR from a patient's serum. This TSH-binding inhibitory immunoglobulins (TBII) assay does not provide information regarding the functionality of TSHR-Ab. TBII assays are cheaper and have good precision. The first generation TBII assays had a sensitivity of 75-95% in untreated Graves' disease patients. Most recently a monoclonal human antibody to TSHR is used. This second generation radio-receptor assay has a sensitivity of 99% with a very high specificity.²¹

3.2.2. *In vitro* bioassays

In vitro bioassays are based on the ability of a patient's serum to stimulate adenylate cyclase and produce cAMP from cultured hamster ovary cells transfected with human TSHR (CHO-R),²³ or rat thyroid cell strain (FRTL-5)²⁴, or human thyroid follicular cells²⁵ are used as a source for functional TSHR. CHO-R system is slightly more sensitive, and requires an easier culture condition than other systems. Advantages of bioassays are that it gives information about the functional property of TSHR-Ab, but bioassays are more expensive, not widely available, have poor precision and sensitivity of more than 90%. These problems have been solved in newer bioassays.

3.2.3. Cellular immunity

The thyroid gland in Graves' disease is characterized by non-homogenous lymphocytic infiltration. Majority of the intrathyroidal lymphocytes are T lymphocytes. B-lymphocytes are much less common than Hashimoto's thyroiditis. Cytokine profile produced by intrathyroidal T lymphocytes suggested that both TH1 & TH2 cells are present in thyroid. The majority of the T-lymphocytes are of TH1 subtype.^{26,27}

TH1 cells are mainly involved in delayed type of hypersensitivity reactions, that produces cytokines like TNF-B, IFN- γ , IL-2, IL-10 and IL-17. TH1 cells are implicated in the pathogenesis of organ specific autoimmune diseases that are mediated mainly by TNF- β subtype, which uniquely produces IL-17. TH 2 cells are mainly responsible for humoral immune responses and they produce cytokines like IL-4, IL-5, IL-6 and IL-13. TH1 cells may also induce antibody formation through secretion of IL-10²⁸. IgG1 subclass of antibodies are selectively induced by TH1 cells. Most of the intrathyroidal T cells are of memory (CD4⁺, CD29⁺) subtype. Concentration of cytotoxic T cells (CD8⁺) are much less in Graves' disease

patients than patients of Hashimoto's thyroiditis. So the functional role of T cells in Graves' disease is primarily that of a helper than a suppressor or cytotoxic role.

4. Pathogenic mechanisms

4.1. Molecular mimicry or specificity crossover

Structural or conformational similarity between different antigens like infectious agents with a self antigen can lead to crossover of specificity or molecular mimicry. Molecular mimicry has been reported between Reoviral antigen and a tissue antigen expressed in multiple endocrine tissues, *Yersinia enterocolitica* and TSHR, Retroviral sequences and TSHR & *Borrelia* and TSHR.

4.2. Bystander effect

There is evidence that bystander activation of local resident antigen specific and nonspecific T- cells by a local viral infection would induce an inflammatory reaction and stimulate the production of cytokines inducing autoimmunity. This bystander activation can also occur with any infection and antigens unrelated to the thyroid gland.

4.3. Aberrant expression of class II HLA Antigens

MHC class II molecules (HLA-DP, DQ, and DR) are not expressed on the normal thyroid epithelial cells but they are expressed on thyroid epithelial cells in patients of autoimmune thyroid disease. This aberrant expression of class II HLA antigens on thyroid epithelia cells can be induced by local thyroid insult which causes production of interferon γ and other cytokines. Interferon γ is able to over express HLA class I molecule and induce the expression of class II molecule on thyroid epithelial cells.

4.4. Cryptic antigens

Autoimmunity results from the loss of tolerance or the ability to differentiate between self and non self. Tolerance induction is a staged process that initiates in the thymus during T-cell maturation. This process depends in part on the presence of peripheral antigens in the thymus. Peripheral antigens are antigens normally expressed in tissues outside of the immune system which are expressed at low levels in thymus. T cells that react strongly to these peripheral molecules in the context of MHC are deleted in thymus. T cells that react with peripheral antigens that are not expressed in the thymus have a greater opportunity to escape tolerance.

4.5. Hygiene hypothesis

Hygiene hypothesis implies that infection may protect from autoimmune diseases rather than precipitating it. Decreased exposure to antigens due to improved living standards can lead to increased risk of autoimmune disorders.

4.6. Super antigens

Super antigens are endogenous or exogenous proteins such as microbial proteins, capable of stimulating a strong immune response through molecular interactions with non-variant parts of the T- cell repertoire and the HLA class II proteins.

5. Pathology

Grossly the thyroid gland is diffusely enlarged with a smooth and hyperemic surface. Rarely the gland is grossly nodular. Consistency of the gland varies from soft to firm. The pyramidal lobe is often prominent.

Microscopically both hypertrophy and hyperplasia are seen. Follicles are small with scanty colloid, and lined by hyperplastic columnar epithelium which can give a pseudopapillary appearance. Vascularity of the gland is increased. There is a varying degree of infiltration by lymphocytes and plasma cells. T cells predominate in the interstitium, whereas B cells and plasma cells predominate in lymphoid follicles.

On electron microscopy there is increased golgi reticulum and mitochondria, and it is also characterized by presence of prominent microvilli.

After treatment with antithyroid drugs and radioiodine, the vascularity of the gland decreases, follicles are enlarged and filled with colloid, and papillary projection regresses.

6. Clinical features

Graves' disease is the most common cause of thyrotoxicosis. Most common age of onset is third to fifth decade of life but it can occur in children and elderly. The hallmark of Graves' disease are signs and symptoms of thyrotoxicosis along with diffuse goiter and typical Graves' orbitopathy. Most of the signs and symptoms are similar to other causes of thyrotoxicosis, but some of the signs and symptoms like orbitopathy, dermopathy or pretibial myxedema and thyroid acropachy are unique to Graves' disease.

Onset of Graves' disease is usually gradual. Signs and symptoms are presents months before the diagnosis, and usually patients do not remember the exact date of onset of symptoms. Onset can be abrupt in some cases. The signs and symptoms are usually more severe than other causes of thyrotoxicosis.

6.1. Thyroid gland

The thyroid gland is diffusely enlarged in Graves' disease, but it can be nodular especially in areas of iodine deficiency where nodular goiter preexists before the onset of Graves' disease. Goiter size is variable; it can range from a normal size thyroid gland to a massively enlarged thyroid. Usually size the of a goiter is two to three times that of normal. A normal size thyroid gland can be seen in as many as 20% of the patients and most of them are elderly. The consistency of the goiter varies from soft to firm but softer than the goiter seen in Hashimoto's

thyroiditis. Thrill and bruit (abn. sounds) can be present over a goiter in severe cases which is due to increased vascularity of the gland. Thrill and bruit are present usually on the upper or lower range of the voice, but sometimes can present only in systole (normal tone). Large goiters with intrathoracic extension can be associated with facial swelling and flushing, and neck vein distension upon raising the arm above the head. This is known as Pemberton sign.

6.2. Graves' ophthalmopathy

Graves' ophthalmopathy is one of the distinctive manifestations of Graves' disease. Overall clinical ophthalmopathy is present in about 50% of Graves' disease patients,²⁹ however CT or MRI reveals extraocular muscle enlargement in about 70% of patients without overt clinical ophthalmopathy.^{30,31} Bimodal age distribution was noted for ophthalmopathy in both men and women. Peak age is 40 to 44 years and 60 to 64 years in women, whereas in men it is 45 to 49 years and 65 to 69 years.

6.2.1. Risk factors for Graves' ophthalmopathy

There is no specific genetic predisposition for graves' ophthalmopathy. Environmental factors are more important for ophthalmopathy.

Smoking

Smoking is a major risk factor for ophthalmopathy. Smoking also increases the risk for worsening of ophthalmopathy after radioiodine treatment. Possible contributors are orbital hypoxia and free radicals present in smoke.^{32,33}

Gender

Graves' disease is predominantly a disease of females (F:M = 8-10:1). In comparison to Graves' disease, ophthalmopathy is relatively more common in males (F:M = 1:1.8-2.8) than females.^{34,35}

Radioiodine

Graves' disease patients treated with radioiodine are at increased risk for onset and worsening of eye disease, as compared to antithyroid drugs alone.^{36,37} This risk can be decreased by concurrent use of corticosteroids.^{38,39}

6.2.2. Pathogenesis

Current evidence supports an autoimmune pathogenesis with important environmental influences, particularly smoking. Orbital muscles, connective tissues, and adipose tissues are infiltrated by lymphocytes and macrophages. TH1 mediated immune response predominates in early stages of disease while TH2 response predominates in late stage.⁴⁰ In response to cytokines secreted by the infiltrating immune cells, orbital fibroblasts start synthesizing and secreting hydrophilic glycosaminoglycans, resulting in edema of orbital

tissues. Additionally adipocytes present in orbit become active and results in expansion of orbital adipose tissues. Both these factors are responsible for expansion of orbital tissues.

6.2.3. *Natural history*

Onset of eye disease usually coincides with that of thyrotoxicosis in 40% of cases, follow it in 40%, and precedes it in 20%.^{41,42} Even when the onset of the two disorders does not coincide, each occurs within 18 months from the onset of the first manifestation. Eye disease usually shows a progressive deterioration lasting for several months followed by a phase of spontaneous improvement lasting up to a year or longer and quiescent stage when inflammatory signs disappear and clinical features stabilizes.

6.2.4. *Signs and symptoms*

The earliest manifestations of ophthalmopathy are usually a sensation of grittiness, eye discomfort, and excess tearing. Other common symptoms are spontaneous retroorbital pain, pain on ocular movement and diplopia. Diplopia is most common in upgaze or in extremes of lateral gaze, because of the involvement of inferior or lateral rectus muscle. About one third of patients have proptosis which can best be detected by visualization of the sclera between the lower border of iris and the lower eyelid when the eyes are kept in primary position.

In severe cases proptosis can cause exposure keratitis and corneal ulcerations. Proptosis is frequently assymetrical. Retraction of the upper eyelid and less commonly of lower eyelid results in lid lag, globe lag, and lagophthalmos. Movements of the lids are jerky, and also there is tremor of lightly closed eyes. These are the non specific manifestations of thyrotoxicosis. Other signs and symptoms are photophobia, swelling of the eyelids, blurring of vision, conjunctival injection and chemosis, periorbital edema.

Decreased visual acuity and color vision, corneal ulceration, and subluxation of globe are present in most severe cases. Blindness may result from corneal ulceration and compression of the optic nerve at the orbital apex due to increased orbital pressure and venous congestion.

Graves' ophthalmopathy can occur in absence of Graves' disease in 10% cases, that's why it is also known as thyroid associated ophthalmopathy, or euthyroid Graves' ophthalmopathy. Graves' ophthalmopathy is usually bilateral but it can be unilateral in upto 10% cases.

6.2.5. *Staging severity of ophthalmopathy*

The acronym NO SPECS was given by American Thyroid Association for severity of ophthalmopathy, where no indicates absence or mild ophthalmopathy, and SPECS indicates more severe degree of involvement, but NO SPECS scheme is inadequate and patients do not necessarily progress from one class to other.

Classes	Ocular symptoms and signs
0	No signs and symptoms
1	Only signs (lid retraction, lid lag, proptosis upto 22 mm)
2	Soft tissue involvement (periorbital edema)
3	Proptosis (>22 mm)
4	Extraocular muscle involvement (diplopia)
5	Corneal involvement
6	Sight loss (optic nerve involvement)

Table 3.

6.2.6. Clinical activity of ophthalmopathy

To know the clinical activity of ophthalmopathy is important, because active disease is more likely to respond to immunosuppressive therapy. Clinical Activity Score (CAS) is used to know the clinical activity. Seven parameters are used in the clinical activity scoring which include spontaneous retrobulbar pain, pain on eye movement, eyelid erythema, conjunctival injection, swelling of the eyelids, inflammation of the caruncle and conjunctival edema or chemosis. Each parameter is assigned 1 point. CAS of more than or equal to 3/7 indicates active ophthalmopathy.

6.3. Thyroid dermopathy

Thyroid dermopathy presents in less than 5% of patients with Graves' disease. It is almost always accompanied by moderate to severe ophthalmopathy. Most commonly it is present over anterior and lateral aspects of leg, hence it is also known as pretibial myxoedema. Less commonly it can present over dorsa of the feet, dorsa of the hands, forearm, face and elbows, particularly after trauma. The typical lesion is a noninflamed, indurated plaque with a deep pink or purple color and an orange skin appearance. Nodular form is the intermediate while elephantiasis is the most severe form of thyroid dermopathy. Thyroid dermopathy occurs due to accumulation of glycosaminoglycans in the dermis and subcutaneous tissues.

6.4. Thyroid acropachy

Thyroid acropachy is the least common manifestation of Graves' disease. It is a form of clubbing, and presents in less than 1 percent of patients of Graves' disease. It is almost always associated with the severe and long standing ophthalmopathy and dermopathy. An alternate diagnosis should be considered in the absence of ophthalmopathy and dermopathy. Deposition of glycosaminoglycans in skin is responsible for thyroid acropachy.

7. Laboratory diagnosis

Diagnosis of Graves' disease can be confirmed by measurement of serum TSH and total thyroxine (TT₄) and triiodothyronine (TT₃). Serum TSH level is suppressed or undetectable with increased TT₄ and TT₃ level in patients of Graves' disease. Serum TSH level is the most sensitive test. The free T₄ (FT₄) and free T₃ (FT₃) levels are increased more than that of TT₄ and TT₃. Measurement of FT₄ and FT₃ are expensive, and there is more chance of laboratory errors. FT₄ and FT₃ can be measured in conditions associated with high serum TBG level like pregnancy, oral contraceptive use and chronic liver disease. In patients of Graves' disease serum T₃ level is proportionately more elevated than the serum T₄ level. In up to 12% of patients, especially in the iodine deficient areas, only TT₃ or FT₃ is elevated with a normal TT₄ or FT₄ level, a condition known as T₃ toxicosis. Conversely in some patients (iodine induced hyperthyroidism, drugs like amiodarone and propranolol which block the conversion of T₄ to T₃), only TT₄ or FT₄ is elevated with normal TT₃ and FT₃ (T₄ toxicosis). Serum thyroglobulin level is high in all cases of thyrotoxicosis except factitious thyrotoxicosis.

Anti TPO antibody can be detected in upto 90%^{43,44} of patients with Graves' disease whereas anti-thyroglobulin antibody is present in 50-80% cases.^{45,46} They are useful in confirming the presence of thyroid autoimmunity but they are of limited diagnostic value. TSHR-Ab assay is very sensitive and specific (upto 98%) for the diagnosis of Graves' disease. But TSHR-Ab assay is quite expensive and not widely available. TSHR-Ab assay is indicated only when clinical and laboratory diagnosis are not clear. Indications for TSHR-Ab assay are:

- Euthyroid Graves' disease, especially when it is unilateral
- Pregnant women with Graves' disease to predict the likelihood of neonatal thyrotoxicosis
- Nodular variant of Graves' disease

TSHR-Ab assay is also a useful indicator of the degree of disease activity. It can also predict the prognosis of Graves' disease. There is more chance of relapse in patients with persistently high TSHR-Ab level after cessation of antithyroid drugs.⁴⁷

Associated hematological abnormalities include increased RBC mass, leukopenia with relative lymphocytosis, monocytosis and eosinophilia, increased factor VIII level. Other associated abnormalities include elevated liver enzymes, bilirubin and ferritin.

7.1. Radioactive iodine uptake (RAIU)

RAIU is not required in each and every case of Graves' disease, but it is useful in excluding thyrotoxicosis caused by thyroiditis, factitious thyrotoxicosis, and type II amiodarone induces thyrotoxicosis. RAIU is absolutely contraindicated in pregnancy. In Graves' disease RAIU is diffusely increased.

7.2. Thyroid ultrasound

In Graves' disease thyroid tissue typically becomes hypoechoic because of reduction in colloid content, increase in thyroid vascularity and lymphocytic infiltration. In colour flow doppler there is a distinct pattern characterized by markedly increased signals, inferior thyroid artery and intrathyroidal artery velocities more than 40 cm/s. This pattern, in conjunction with a hypoechoic pattern allows distinction from Hashimoto's thyroiditis.

7.3. Differential diagnosis

If a patient presented with diffuse goiter with clinical and biochemical thyrotoxicosis, ophthalmopathy and positive autoimmune markers like anti TPO antibody or TSHR-Ab, diagnosis of Graves' disease is straight forward. In absence of these classical features radionuclide scan (I^{123} , I^{131} , Tc^{99m}) is the most reliable distinguishing test. In case of Graves' disease there is diffuse and high uptake, whereas in patients with toxic adenoma or toxic multinodular goiter there is patchy uptake. In patients with thyroiditis and factitious thyrotoxicosis there is decreased uptake. In patients with TSH producing adenoma, there is also a diffuse goiter, but TSH is inappropriately normal or increased instead of suppress TSH of graves' disease. Panic attacks, mania, pheochromocytoma and malignancy can be easily ruled out by thyroid function test.

8. Treatment

There is no ideal treatment option present targeting the basic pathogenic mechanisms of Graves' disease. Available treatment options target the increased synthesis of thyroid hormones by antithyroid drugs, or ablation of thyroid tissue by surgery or radioiodine. Antithyroid drugs are the predominant therapy in Europe and Japan whereas radioiodine is first line of treatment in USA. No single treatment is optimal and patients often require multiple treatment.

8.1. Antithyroid drugs

Thionamides are the main antithyroid drugs which includes propylthiouracil, carbimazole and methimazole. Thionamides act by inhibiting the enzyme thyroid peroxidase, reducing the oxidation and organification of iodide and coupling of iodotyrosines. Carbimazole is not an active drug, and in body it is converted to active metabolite methimazole. Propylthiouracil, in addition to inhibiting thyroid hormone synthesis, also inhibits the peripheral conversion of T_4 to T_3 . Methimazole is ten times more potent than propylthiouracil. The half life of methimazole is about 6 hours while that of propylthiouracil is about 90 minutes. Duration of action of methimazole is more than 24 hours while that of propylthiouracil is 12-24 hours. Transplacental transfer of propylthiouracil is low. Antithyroid drugs do not block the release of preformed hormones, so euthyroidism is not obtained until intrathyroidal hormone store is depleted. These drugs also reduce thyroid antibody level.

Antithyroid drugs can be used as a primary treatment or as a preparatory treatment before radioiodine or surgery. The antithyroid drugs are usually started in higher doses. The starting dose of methiomazole or carbimazole is 10-20 mg every 8-12 hours and that of propylthiouracil is 100-200 mg every 6-8 hours. Once euthyroidism is achieved which usually takes 4-5 weeks, methimazole can be given in single daily dose while propylthiouracil is given in multiple daily doses throughout the treatment. There are two treatment strategies for using antithyroid drugs. In titration regimen, antithyroid drugs are started in high doses and doses can be gradually decreased to maintain euthyroid state. In Block and Replace regimen, the antithyroid drugs are maintained in high doses and subsequently levothyroxine is added to maintain the euthyroid status. At present there is no proven advantage of block and replace regimen over the titration regimen.

Patients should be reviewed clinically and biochemically after every 3-4 weeks. Doses of antithyroid drugs are adjusted based on the TT_4 or FT_4 level, as TSH level often remains suppressed for several months. The usual daily maintenance dose of carbimazole or methimazole is 2.5 – 10 mg and that of propylthiouracil is 50-100 mg. When the TSH level becomes normal, it can also be used to monitor therapy. The size of the goiter decreases in about 30-50% of patients during treatment. In remaining patients it may remain unchanged or even enlarge. Increase in goiter size is one of the earliest manifestations of iatrogenic hypothyroidism. The other features are weight gain, lethargy, fatigue and other signs of mild hypothyroidism.

Maximum remission rate with antithyroid drugs is 30-50%, which can be achieved by continuation of the drug for 6-18 months or even longer. Most of the relapses occur within first 3-6 months after discontinuation of drug. Suppressed TSH level below the normal limit is the first signal of relapse even in the presence of a normal serum T_4 level. In most of the studies, the relapse rate is 50-80%.^{48,49} The most important predictor of relapse is goiter size.⁵⁰ Other factors influencing recurrence of Graves' disease include high TSHR-Ab concentration, large iodine intake, marked residual goiter, short duration of antithyroid drug treatment and previous recurrence. Factors which favor long term remission after therapy include the initial presence of T_3 toxicosis, a small goiter, decrease in the size of goiter, and return of TSH to normal during treatment, the return of serum thyroglobulin to normal, and low iodine diet.

Most common side effects of thionamides are pruritus, skin rash, urticaria, fever and arthralgia. These may resolve spontaneously or after substituting another drug. Rare but major side effects are hepatitis, cholestasis, SLE like syndrome, ANCA positive vasculitis and most importantly agranulocytosis. Major side effects occur in less than 1% of patients. Antithyroid drugs should be stopped and not restarted if patients develop major side effects.

8.2. Iodine and iodine containing components

Inorganic iodine acts in many ways in thyrotoxicosis. Iodine blocks its own transport in thyroid, inhibits iodine organification and inhibits the release of hormones. Inhibition of iodine organification by inorganic iodide is known as Wolf-Chaikoff effect. The major action of

iodine is inhibition of hormone release. Iodine also decreases the vascularity of the thyroid gland. All of these effects of iodine are transient and last only for a few days or weeks. Today iodine is used only for preoperative preparation for Graves' disease and in the management of thyrotoxic crisis. The usual dose of Lugol's solution is 3-5 drops three times per day and that of SSKI is 2-3 drops twice daily. Iodine decreases the effect of subsequently administered thionamides and radioiodine for several weeks.

Iodinated radio contrast agents like iopanoic acid, and sodium ipodate act by blocking the peripheral conversion of T4 to T3 and inhibition of hormone release. They are ideally used in emergency situations when rapid control of thyrotoxicosis is needed or in preoperative preparation or while awaiting the response of radioiodine.

8.3. Thiocyanate and perchlorate

They act by initiating the transport of iodine to thyroid gland.

8.4. Lithium

Lithium also acts as a thyroid constipating agent (block the release of thyroid hormones). Lithium also potentiates the beneficial effect of radioiodine. The usual dose of lithium is 450-900 mg per day in divided doses. Serum lithium concentration should be maintained at 1mEq/L. No adverse effects are reported with this dose of lithium.

8.5. Glucocorticoids

Dexamethasone in high doses (8mg/day) decreases the T4 secretion by the thyroid gland, inhibits the peripheral conversion of T4 to T3, and has an immunosuppressive effect. The effect on peripheral conversion of T4 to T3 is additive to propylthiouracil. Glucocorticoids are indicated for the treatment of ophthalmopathy, dermopathy and in thyrotoxic crisis.

8.6. β - blockers

β - blockers do not affect the synthesis or secretion of thyroid hormones. Many of the symptoms & signs of thyrotoxicosis are due to hypersensitivity of the sympathetic nervous system to thyroid hormones. Thus use of β - blockers in thyrotoxicosis, improve the signs and symptoms mediated by the sympathetic nervous system. Tachycardia, palpitation, tremor, anxiety, excess sweating, lid retraction improves with β - blockers. Propranolol has additional advantage over other β - blockers by inhibiting the peripheral conversion of T4 to T3. β - blockers reduce cardiac output without altering oxygen consumption, but can have adverse effects in liver, where the arteriovenous oxygen difference is already elevated in the hyperthyroid state. Propranolol is the most commonly used agent but other β - blockers can also be used. It is used in a dose of 20-60 mg every 6-8 hours. Short acting agents like esmolol is used for intravenous purpose. Long acting agents like atenolol or metoprolol are used for prolonged treatment. β - blockers should be rapidly tapered and discontinued once stable euthyroidism is achieved.

8.7. Radioiodine

Radioiodine is one of the first line therapies for the Graves' disease. Among different isotopes of radioiodine, I^{131} is the agent of choice. I^{131} is a β -emitter isotope. After oral administration, I^{131} is completely absorbed and rapidly concentrated in thyroid follicular cells. β - particles, which are emitted by I^{131} , destroy the thyroid follicular cells that results in reduced thyroid hormone synthesis. Initially destruction of thyroid follicular cells results in release of preformed hormones that can precipitate the thyrotoxic crisis. Weeks to months are required for control of thyrotoxicosis. Long term effects of radioiodine include atrophy and fibrosis, and a chronic inflammatory response resembling Hashimoto's thyroiditis.

Radioiodine is given as single oral dose. Three outcomes of radioiodine treatment are possible- patients become euthyroid, remain thyrotoxic, or become permanently hypothyroid. The dose of radioiodine depends on the size of gland, the uptake of I^{131} and its subsequent rate of release. The dose of radioiodine ranges from 80-200 $\mu\text{Ci/gm}$ of thyroid tissues. A total dose of 20 mCi achieves thyroid ablation in almost all patients and results in permanent hypothyroidisms in 75-90% of patients.⁵¹ The incidence of post radioiodine hypothyroidism in first year is 25% and steadily increases thereafter at a rate of 5% per year. When required, the second dose of radioiodine should be given at least 6 months after the first dose. Failure of radioiodine treatment is more common in patients with large goiter, rapid iodine turnover and adjunctive antithyroid drugs too soon after radioiodine. Prior use of antithyroid drugs decreases the risk of thyrotoxic crisis. Chance of worsening ophthalmopathy can also be reduced by antithyroid drugs. Antithyroid drugs should be stopped 3 to 8 days prior to radioiodine treatment and should be restarted after 7 days when required. Propylthiouracil may cause radio-resistance, but it not a major concern. Short term side effects of radioiodine include transient exacerbation of thyrotoxicosis in the first few months, transient worsening of ophthalmopathy, acute radiation thyroiditis in the first week. Radiation thyroiditis may lead to transient worsening of thyrotoxicosis and ophthalmopathy. Presence of mild to moderate ophthalmopathy is not a contraindication for radioiodine treatment. Concomitant use of oral glucocorticoids, decreases the risk of worsening ophthalmopathy. Long term side effect of radioiodine is permanent hypothyroidism. Initially there was a concern regarding possible carcinogenic effect and risk of genetic damage after radioiodine treatment, but now it is proven that there is no association between radioiodine treatment and thyroid carcinoma, leukemia, solid tumors and genetic damage.⁵² Thyroid cancer is associated with low dose of I^{131} rather than higher dose of I^{131} in children.⁵³ Some centers use radioiodine even in children of 10 years of age or younger, but still there is no consensus regarding use of radioiodine for persons younger than 16 to 18 years.

8.8. Surgery

Surgery is a form of ablative therapy. Enough thyroid tissue is removed by surgery to reduce the synthesis of thyroid hormones and prevent recurrence. Two type of thyroid surgery are used for Graves' disease. In subtotal thyroidectomy, the bulk of the gland is removed and only about 2 gm (0.5%) of thyroid tissue is left in both lobes. In near total thyroidectomy, most

of the thyroid gland is removed, with only subcentimeter fragments left around recurrent laryngeal nerve and parathyroid glands. Subtotal thyroidectomy was the procedure of choice in the past, but it is associated with higher recurrence rate. Now, near total thyroidectomy is used most often. Near total thyroidectomy is associated with a more permanent chance of hypothyroidism but less chance of recurrence than with subtotal thyroidectomy.

Complications of thyroid surgery depends on the skill and experience of the operating surgeon. In specialized hands, the complication rate is as low as 2%, whereas complication rates increase up to 10-15% in non specialized centers. Post Operative bleeding is the most serious complication and can be fatal by producing asphyxia if it is not evacuated immediately. Other complications such as thyroid storm, injury to recurrent laryngeal nerve and hypoparathyroidism are specific to thyroid surgery. Thyroid storm is typically rare. Injury to recurrent laryngeal nerves causes dysphonia that usually improves with time, but that may leave the patient slightly hoarse. Hypoparathyroidism can be transient or permanent. Transient hypoparathyroidism is due to removal of the some parathyroid and impairment of blood supply to parathyroid glands, whereas permanent hypoparathyroidism is due to inadvertent removal of all 4 glands. Transient hypoparathyroidism usually occurs on day 1-7 postoperatively. Severe symptomatic hypoparathyroidism should be treated by intravenous calcium gluconate. Oral calcium (upto 3 gm/day) is sufficient for milder cases. Immediate postoperative hypocalcemia is due to 'hungry bone syndrome'. Recurrence of hyperthyroidism and permanent hypothyroidism are inversely related and depends on the amount of thyroid tissue left. In case of recurrence, radioiodine should be used as treatment, as a second surgery is technically difficult.

Preoperative preparation

Preoperative use of thionamides is associated with a lower morbidity and mortality. Preoperative thionamides are recommended to achieve euthyroidism and to deplete the hormone store. Preoperative use of inorganic iodine decreases the gland size and vascularity. In case of emergency surgery, oral cholecystographic agents are the fastest way to obtain euthyroidism. The goal of preoperative management is to maintain euthyroid states by thionamides and then to induce involution of the gland by the inorganic iodine. β blockers can be used in preoperative preparation.

Choice of therapy

Choice of therapy depends on the patient preference, personal experience of the treating doctor and availability of the treatment options. All three treatment options (antithyroid drugs, radioiodine, surgery) can be used as first line therapy. In most of Europe, antithyroid drugs are the preferred treatment whereas in USA, radioiodine is the preferred treatment. Primary choice for treatment in children and young adults up to 18 years of age is antithyroid drugs, although radioiodine is not associated with any adverse events. Pregnancy should be delayed for 6-12 months after radioiodine treatment. Presence of severe ophthalmopathy is a contraindication for radioiodine treatment. Surgery is the preferred treatment for patients with large goiter, especially if compressive symptoms are present, endemic goiter with multiple cold nodules, and suspected malignancy.

Thyroid ophthalmopathy

Mild to moderate ophthalmopathy does not require any specific treatment. General measures include control of thyrotoxicosis, smoking cessation, dark glasses with side frame for photophobia and sensitivity to air, artificial tears (1% methyl cellulose) or eye ointment for eye discomfort and dry eye, eye patches or taping during sleep for lagophthalmos, elevation of the head end for periorbital edema, prism for correction of mild diplopia.

Other patients with more severe signs and symptoms affecting daily lives to a significant extent, may benefit from immunosuppressive therapy in active disease or surgical decompression in case of inactive disease. Severe ophthalmopathy with optic neuropathy and corneal ulcer is considered an emergency.

Glucocorticoids

Oral glucocorticoids is initiated at a relatively high dose, such as 40-80 mg of prednisolone per day. After 2-4 weeks, the daily dose is tapered by 2.5-10 mg every 2-4 weeks. Improvement in soft tissue inflammation begins within 1-2 days. Intravenous methylprednisolone pulse therapy is more effective and better tolerated than oral prednisolone.⁵⁴ 500 mg of methylprednisolone per week for 6 weeks followed by 250 mg of methylprednisolone per week for 6 weeks is most commonly used regimen. Cyclosporine can also be used either as a single therapy or in combination with oral prednisolone. Combination therapy of cyclosporine with prednisolone is more effective than either drug alone.⁵⁵

Orbital Radiotherapy

Orbital radiotherapy is well tolerated and provides benefits in approximately two thirds of the patients. This treatment is steroid sparing rather than steroid replacing therapy.

Other immunomodulatory therapy

Rituximab, azathioprine, cyclophosphamide, ciamexon, pentoxifylline and intravenous immunoglobulins have some benefit and are currently under trial.

Orbital decompression

Indications for orbital decompression include optic neuropathy, severe proptosis, vision threatening ocular exposure, debilitating retrobulbar and periorbital pain and intolerable corticosteroid side effects. Transantral orbital decompression with removal of a portion of medial wall and the orbital floor is most commonly used procedure. Up to a 5 mm reduction in proptosis can be achieved by orbital decompression. Orbital decompression can cause onset or worsening of diplopia.

Thyroid storm

Thyroid storm or thyrotoxic crisis is a life threatening exacerbation of hyperthyroidism. Most of the cases of thyroid storm are associated with Graves' disease, but it can also occur with toxic multinodular goitre. Precipitating factors for thyroid storm include infection,

trauma, thyroid surgery, radioiodine, diabetic ketoacidosis, stroke etc. It can present with fever, tachycardia, arrhythmias, profuse sweating, diarrhea and vomiting, confusion, delirium, seizures, jaundice, coma, congestive heart failure, hypotension. Thyroid storm is associated with very high mortality rate of up to 30%, even with treatment.

Treatment of thyroid storm requires strict monitoring and proper care. Precipitating factors should be identified and treated. Supportive treatments include cooling blankets and drugs like acetaminophen, chlorpromazine or meperidine for hyperthermia, oxygen inhalation, and intravenous fluids. The antithyroid drug of choice is propylthiouracil, but carbimazole can also be used. Propylthiouracil 600 mg is given as a loading dose by mouth or nasogastric tube, or per rectum followed by 200-300 mg every 6-8 hourly. One hour after the first dose of propylthiouracil, stable iodide is given in the form of SSKI (3 drops twice daily) or Lugol's iodine (10 drops twice daily). Propranolol should also be given in a dose of 40-80 mg orally every 6 hours or 2 mg intravenously every 4 hours. If β blockers are contraindicated, calcium channel blockers like diltiazem can be used to control tachycardia. Large doses of dexamethasone (8 mg) by oral or intravenous route should be given to block the release of hormone from gland and peripheral conversion of T4 to T3.

Author details

N.K. Agrawal and Ved Prakash

*Department of Endocrinology and Metabolism, Institute of Medical Sciences,
Banaras Hindu University, Varanasi, India*

Manuj Sharma

Department of Medicine, Gandhi Medical College, Bhopal

9. References

- [1] Graves RJ: Newly observed affection of the thyroid. *London Med Surg J* 1835; 7:515-523.
- [2] Parry CH: *Collections from the Unpublished Medical Writings of the Late Caleb Hillier Parry.* London, Underwood, 1825.
- [3] von Basedow KA: Exophthalmos durch hypertrophie des zellgewebes in der Augenhole. *Wochenschr Ges Heilk Berl* 1840; 6:197.
- [4] Laurberg P, Pedersen KM, Vestergaard H, et al: High incidence of multinodular toxic goitre in the elderly population in a low iodine intake area vs. high incidence of Graves' disease in the young in a high iodine intake area: comparative surveys of thyrotoxicosis epidemiology in East-Jutland Denmark and Iceland. *J Intern Med* 1991; 229:415-420.
- [5] Furszyfer J, Kurland LT, McConahey WM, et al: Epidemiologic aspects of Hashimoto's thyroiditis and Graves' disease in Rochester Minnesota (1935-1967), with special reference to temporal trends. *Metabolism* 1972; 21:197-204.
- [6] Aghini-Lombardi F, Antonangeli L, Martino E, et al: The spectrum of thyroid disorders in an iodine-deficient community: the Pescopagano survey. *J Clin Endocrinol Metab* 1999; 84:561-566

- [7] Aghini-Lombardi F, Antonangeli L, Martino E, et al: The spectrum of thyroid disorders in an iodine-deficient community: the Pescopagano survey. *J Clin Endocrinol Metab* 1999; 84:561-566
- [8] Bülow Pedersen I, Laurberg P, Knudsen N, et al: Increase in incidence of hyperthyroidism predominantly occurs in young people after iodine fortification of salt in Denmark. *J Clin Endocrinol Metab* 2006; 91:3830-3834,2006.
- [9] Chiovato L, Lapi P, Fiore E, et al: Thyroid autoimmunity and female gender. *J Endocrinol Invest* 1993; 16:384-391.
- [10] Amino N, Tada H, Hidaka Y: Postpartum autoimmune thyroid syndrome: a model of aggravation of autoimmune disease. *Thyroid* 1999; 9:705-713.
- [11] Bartalena L: Smoking and Graves' disease. *J Endocrinol Invest* 2002; 25:402.
- [12] Holm IA, Manson JE, Michels KB, et al: Smoking and other lifestyle factors and the risk of Graves' hyperthyroidism. *Arch Intern Med* 2005; 165:1606-1611.
- [13] Adams DD, Purves HD: Abnormal responses in the assay of thyrotropin. *Proc Univ Otago Med Sch* 1956; 34:11-12.
- [14] McKenzie JM, Zakarija M: Fetal and neonatal hyperthyroidism and hypothyroidism due to maternal TSH receptor antibodies. *Thyroid* 1992; 2:155-159.
- [15] McLachlan SM, Dickinson AM, Malcolm A, et al: Thyroid autoantibody synthesis by cultures of thyroid and peripheral blood lymphocytes. I: Lymphocyte markers and response to pokeweed mitogen. *Isr J Med Sci* 1983; 52:45-53.
- [16] Okuda J, Akamizu T, Sugawa H, et al: Preparation and characterization of monoclonal antithyrotropin receptor antibodies obtained from peripheral lymphocytes of hypothyroid patients with primary myxedema. *J Clin Endocrinol Metab* 1994; 79:1600-1604.
- [17] Bolton J, Sanders J, Oda Y, et al: Measurement of thyroid stimulating hormone receptor autoantibodies by ELISA. *Clin Chem* 1999; 45:2285-2287.
- [18] Costagliola S, Morgenthaler NG, Hoermann R, et al: Second generation assay for thyrotropin receptor antibodies has superior diagnostic sensitivity for Graves disease. *J Clin Endocrinol Metab* 1999; 84:90-97.
- [19] Pinchera A, Liberti P, Martino E, et al: Effects of antithyroid therapy on the long-acting thyroid stimulator and the antithyroglobulin antibodies. *J Clin Endocrinol* 1969; 29:231-289.
- [20] Fenzi GF, Hashizume K, Roudeboush CP, et al: Changes in thyroid-stimulating immunoglobulins during antithyroid therapy. *J Clin Endocrinol Metab* 1979; 48:572-576.
- [21] Costagliola S, Morgenthaler NG, Hoermann R, et al: Second generation assay for thyrotropin receptor antibodies has superior diagnostic sensitivity for Graves' disease. *J Clin Endocrinol Metab* 1999; 84:90-97.
- [22] Schott M, Scherbaum WA, Morgenthaler NG: Thyrotropin receptor autoantibodies in Graves' disease. *Trends Endocrinol Metab* 2005; 16:243-248.
- [23] Ludgate M, Perret J, Parmentier M, et al: Use of the recombinant human thyrotropin receptor (TSH-R) expressed in mammalian cell lines to assay TSH-R autoantibodies. *Mol Cell Endocrinol* 1990; 73:R13-R18.

- [24] Vitti P, Valente WA, Ambesi-Impiombato FS, et al: Graves' IgG stimulation of continuously cultured rat thyroid cells: a sensitive and potentially useful clinical assay. *J Endocrinol Invest* 1982; 5:179-182.
- [25] Rapoport B, Greenspan FS, Filetti S, et al: Clinical experience with a human thyroid cell bioassay for thyroid-stimulating immunoglobulin. *J Clin Endocrinol Metab* 1984; 58:332-338.
- [26] Mariotti S, del Prete GF, Mastromauro C, et al: The autoimmune infiltrate of Basedow's disease: analysis of clonal level and comparison with Hashimoto's thyroiditis. *Exp Clin Endocrinol* 1991; 97:139-146.
- [27] Watson PF, Pickerill AP, Davies R, et al: Analysis of cytokine gene expression in Graves' disease and multinodular goiter. *J Clin Endocrinol Metab* 1994; 79:355-360.
- [28] Leovey A, Nagy E, Balazs G, et al: Lymphocytes resided in the thyroid are the main source of TSH-receptor antibodies in Basedow's-Graves' disease?. *Exp Clin Endocrinol* 1992; 99:147-150.
- [29] Burch HB, Wartofsky L: Graves' ophthalmopathy: current concepts regarding pathogenesis and management. *Endocr Rev* 1993; 14:747-793.
- [30] Forbes G, Gorman CA, Brennan MD, et al: Ophthalmopathy of Graves' disease: computerized volume measurements of the orbital fat and muscle. *Am J Neuroradiol* 1986; 7:651-656
- [31] Villadolid MC, Nagataki S, Uetani M, et al: Untreated Graves' disease patients without clinical ophthalmopathy demonstrate a high frequency of extraocular muscle (EOM) enlargement by magnetic resonance. *J Clin Endocrinol Metab* 1995; 80:2830-2833.
- [32] Metcalfe RA, Weetman AP: Stimulation of extraocular muscle fibroblasts by cytokines and hypoxia: possible role in thyroid-associated ophthalmopathy. *Clin Endocrinol (Oxf)* 1994; 40:67-72.
- [33] Burch HB, Lahiri S, Bahn R, et al: Superoxide radical production stimulates human retroocular fibroblast proliferation in Graves' ophthalmopathy. *Exp Eye Res* 1997; 65:311-316.
- [34] Vanderpump MP, Tunbridge WM, French JM, et al: The incidence of thyroid disorders in the community: A twenty-year follow-up of the Whickham survey. *Clin Endocrinol (Oxf)* 1995; 43:55-68.
- [35] Perros P, Kendall-Taylor P: Pathogenetic mechanisms in thyroid-associated ophthalmopathy. *J Intern Med* 1992; 231:205-211.
- [36] Fernandez-Sanchez JR, Vara-Thorbeck R, Garbin-Fuentes I, et al: Graves' ophthalmopathy after subtotal thyroidectomy and radioiodine therapy. *Br J Surg* 1993; 80:1134-1136.
- [37] Bartalena L, Pinchera A, Martino E, et al: Relation between therapy for hyperthyroidism and the course of Graves' ophthalmopathy. *N Engl J Med* 1998; 338:73-78.
- [38] Bartalena L, Marcocci C, Bogazzi F, et al: Use of corticosteroids to prevent progression of Graves' ophthalmopathy after radioiodine therapy for hyperthyroidism. *N Engl J Med* 1989; 321:1349-1352..
- [39] Weetman AP, Wiersinga WM: Current management of thyroid-associated ophthalmopathy in Europe: results of an international survey. *Clin Endocrinol (Oxf)* 1998; 49:21-28.

- [40] Aniszewski JP, Valyasevi RW, Bahn RS: Relationship between disease duration and predominant orbital T cell subset in Graves' ophthalmopathy. *J Clin Endocrinol Metab* 2000; 85:776-780.
- [41] Bartalena L, Pinchera A, Marcocci C: Management of Graves' ophthalmopathy: reality and perspectives. *Endocr Rev* 2000; 21:168-199.
- [42] Bartalena L, Wiersinga WM, Pinchera A: Graves' ophthalmopathy: state of the art and perspectives. *J Endocrinol Invest* 2004; 27:295-301.
- [43] Mariotti S, Caturegli P, Piccolo P, et al: Antithyroid peroxidase autoantibodies in thyroid diseases. *J Clin Endocrinol Metab* 1990; 71:661-669.
- [44] McLachlan SM, Rapoport B: Thyroid peroxidase as an autoantigen. *Thyroid* 2007; 17:939-994.
- [45] McLachlan SM, Rapoport B: Why measure thyroglobulin autoantibodies rather than thyroid peroxidase autoantibodies?. *Thyroid* 2004; 14:510-520.
- [46] Mariotti S, Barbesino G, Caturegli P, et al: Assay of thyroglobulin in serum with thyroglobulin autoantibodies: an unobtainable goal?. *J Clin Endocrinol Metab* 1995; 80:468-472.
- [47] Davis TF: Thyroid stimulating antibodies predict hyperthyroidism. *J Clin Endocrinol Metab* 1998; 83:3777-3781.
- [48] Hedley AJ, Young RE, Jones SJ, et al: Antithyroid drugs in the treatment of hyperthyroidism of Graves' disease: long-term follow-up of 434 patients. Scottish Automated Follow-Up Register Group. *Clin Endocrinol (Oxf)* 1989; 31:209-218.
- [49] Schleusener H, Schwander J, Fischer C, et al: Prospective multicentre study on the prediction of relapse after antithyroid drug treatment in patients with Graves' disease [published erratum appears in *Acta Endocrinol (Copenh)* 1989 Aug; 121(2):304]. *Acta Endocrinol (Copenh)* 1989; 120:689-701.
- [50] Vitti P, Rago T, Chiovato L, et al: Clinical features of patients with Graves' disease undergoing remission after antithyroid drug treatment. *Thyroid* 1997; 7:369-375.
- [51] Razvi S, Basu A, McIntyre EA, et al: Low failure rate of fixed administered activity of 400 MBq of I131 with pre-treatment with carbimazole for thyrotoxicosis: the Gateshead Protocol. *Nucl Med Commun* 2004; 25:675-682.
- [52] Franklyn JA, Daykin J, Drolc Z, et al: Long-term follow-up of treatment of thyrotoxicosis by three different methods. *Clin Endocrinol (Oxf)* 1991; 34:71-76.
- [53] Dobyns BM, Sheline GE, Workman JB, et al: Malignant and benign neoplasms of the thyroid in patients treated for hyperthyroidism: a report of the Cooperative Thyrotoxicosis Therapy Follow-Up Study. *J Clin Endocrinol Metab* 1974; 38:976-998.
- [54] Kahaly GJ, Pitz S, Hommel G, et al: Randomized, single blind trial of intravenous versus oral steroid monotherapy in Graves' orbitopathy. *J Clin Endocrinol Metab* 2005 Sep; 90(9):5234-5240.
- [55] Prummel MF, Mourits MP, Berghout A, et al: Prednisone and cyclosporine in the treatment of severe Graves' ophthalmopathy. *N Engl J Med* 1989; 321:1353-1359.